

FAX FORM TO: 1.877.597.3070

PHONE: 1.800.473.3516

EMAIL: specialty@wellpartner.com

Complete the following or include demographic sheet.

1. PATIENT INFORMATION

Name: _____

Address: _____

City, State, ZIP: _____

Primary Phone: - - - - - DOB: / /

Alternate Phone: - - - - - Gender: _____

Email: _____

Primary Language: _____ Last Four of SSN: _____

2. PRESCRIBER INFORMATION

Name: _____

DEA #: _____ NPI #: _____ State Lic. #: _____

Group or Hospital: _____

Address: _____

City, State, Zip: _____

Phone: - - - - - Fax: - - - - -

Contact Person: _____ Phone: - - - - -

3. INSURANCE INFORMATION *Fax copy of prescription and insurance cards with this form, if available (front and back)*

Primary Insurance Company Name: _____ Secondary Insurance Company Name: _____

Primary Cardholder Name: _____ Secondary Cardholder Name: _____

Relationship: Self Spouse/Partner Child/Dependent Relationship: Self Spouse/Partner Child/Dependent

Phone: - - - - - Member ID: _____ Group #: _____ Phone: - - - - - Member ID: _____ Group #: _____

4. DIAGNOSIS AND CLINICAL INFORMATION

Needs by Date: / / Ship to: Patient Office Other:

Date of Diagnosis: / /

D63.8 Anemia in other chronic diseases
 Other diagnosis (please include ICD-10 code): _____

Allergies: _____

Other Medications:
 Ribavirin (dose/strength: _____)
 Other(s): _____

Is injection training necessary? Yes No
 If no, reason: Patient independent
 MD office trained patient or referred to other trainer

Lab Data (if applicable):
 Last clinic visit: / / HGB: _____ HCT: _____ GFR: _____ Ferritin: _____
 Platelets: _____ Creatinine clearance: _____ mL/min Serum creatinine: _____ mg/dl

Is patient's blood pressure under control? Yes No
 Is patient being monitored for thrombotic/cardiac events? Yes No
 Has anemia occurred due to hemolysis, nutritional deficiencies, or GI bleeds? Yes No
 If patient is currently taking hematopoetics, provide evaluation of response to therapy: _____

HGB rise ≥ _____ g/dL and/or HCT ≥ 3% Yes No
 Dose or administration frequency is adjusted until Hgb level is ≤ 12 g/dL Yes No
 Hgb increased more than 1 g/dL in a two-week period or Hgb level is approaching 12 g/dL Yes No
 Dose to be reduced to avoid rapid rise in Hgb level Yes No

5. PRESCRIPTION INFORMATION

Medication	Dose/Strength	Directions	Qty	Refills
<input type="radio"/> Aranesp™	<input type="radio"/> 25 mcg <input type="radio"/> 100 mcg <input type="radio"/> 300 mcg <input type="radio"/> 40 mcg <input type="radio"/> 150 mcg <input type="radio"/> 500 mcg <input type="radio"/> 60 mcg <input type="radio"/> 200 mcg <input type="radio"/> Autoinjector <input type="radio"/> PFS <input type="radio"/> Vial	<input type="radio"/> Inject the entire contents of autoinjector/syringe SQ once every other week <input type="radio"/> Inject the entire contents of autoinjector/syringe SQ once a week <input type="radio"/> Other:		
<input type="radio"/> Epoprostenol™	<input type="radio"/> 2,000 u/ml (SDV) <input type="radio"/> 10,000 u/ml (SDV) <input type="radio"/> 3,000 u/ml (SDV) <input type="radio"/> 10,000 u/ml 2 ml vial (MDV) <input type="radio"/> 4,000 u/ml (SDV) <input type="radio"/> 20,000 u/ml 1 ml vial (MDV)	<input type="radio"/> Single-dose vial (SDV): Inject entire contents of 1 vial SQ: <input type="radio"/> Once a week <input type="radio"/> 3 times a week <input type="radio"/> Other: <input type="radio"/> Multi-dose vial (MDV): Inject _____ ml (_____ units) SQ: <input type="radio"/> Once a week <input type="radio"/> 3 times a week <input type="radio"/> Other:		
<input type="radio"/> Leukine™	<input type="radio"/> 250 mcg vial (lyophilized) <input type="radio"/> 500 mcg vial (liquid)	<input type="radio"/> Administer _____ mcg once a day for _____ days. (Circle IV or SC) <input type="radio"/> Other:		
<input type="radio"/> Neulasta™	6 mg prefilled syringes (PFS)	<input type="radio"/> Inject 6 mg SQ day after chemotherapy, every _____ days. <input type="radio"/> Other:		
<input type="radio"/> Neumega™	5 mg vial kit	<input type="radio"/> Mix and administer 50 mcg/kg once a day for _____ days. <input type="radio"/> Other:		
<input type="radio"/> Neupogen™	<input type="radio"/> 300 mcg <input type="radio"/> PFS <input type="radio"/> 480 mcg <input type="radio"/> Vial	<input type="radio"/> Administer _____ mcg once a day for _____ days. (Circle IV or SC) <input type="radio"/> Other:		
<input type="radio"/> Procrit™	<input type="radio"/> 2,000 u/ml (SDV) <input type="radio"/> 10,000 u/ml (SDV) <input type="radio"/> 3,000 u/ml (SDV) <input type="radio"/> 10,000 u/ml 2 ml vial (MDV) <input type="radio"/> 4,000 u/ml (SDV) <input type="radio"/> 20,000 u/ml 1 ml vial (MDV)	<input type="radio"/> Single-dose vial (SDV): Inject entire contents of 1 vial SQ: <input type="radio"/> Once a week <input type="radio"/> 3 times a week <input type="radio"/> Other: <input type="radio"/> Multi-dose vial (MDV): Inject _____ ml (_____ units) SQ: <input type="radio"/> Once a week <input type="radio"/> 3 times a week <input type="radio"/> Other:		
<input type="radio"/> Other				

Ancillary supplies and kits will be provided as needed for administration.

6. PRESCRIBER SIGNATURE

X _____ / / X _____ / /

DISPENSE AS WRITTEN DATE PRODUCT SUBSTITUTION PERMITTED DATE

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address or telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by express authority of the sender to the named addressee.